

## **The Compositional Profile of the Enceladus Dust Plume I. The Data**

Frank Postberg<sup>1</sup>, Juergen Schmidt<sup>2</sup>, Sascha Kempf<sup>3</sup>, Jon Hillier<sup>4</sup>, Ralf Srama<sup>5</sup>

<sup>1</sup>*Universitaet Heidelberg, Germany,*

<sup>2</sup>*University of Potsdam, Germany,*

<sup>3</sup>*Max Planck Institut fuer Kernphysik, Heidelberg, Germany,*

<sup>4</sup>*Planetary and Space Sciences Research, The Open University, United Kingdom*

<sup>5</sup>*Max-Planck-Institute for Nuclear Physics, Germany*

Data obtained by the Cassini spacecraft during recent close flybys of Enceladus will be presented. Prior compositional measurements of E ring grains with Cassini's Cosmic Dust Analyser (CDA) suggested salt-rich water as the dominant source of Enceladus' famous plume. Although the E ring dust population is dominated by ice particles stemming from Enceladus it was unclear how representative the E ring particles were of the original plume grain ensemble. During 2008 and 2009 Cassini passed deep into the plumes on several occasions, allowing the CDA team to analyse the compositions of freshly ejected plume particles for the first time. From this information a compositional profile of the plume has been inferred, showing that close to Enceladus' surface its composition differs significantly from that of the E ring. The observations can only be reproduced with a liquid water plume source.