## Quasi-Periodic Pulsations in the Gamma-ray, Hard X-ray and Microwave Emission of a Solar Flare

VALERY M. NAKARIAKOV<sup>1</sup>, CLAIRE FOULLON<sup>1</sup>, IRINA N. MYAGKOVA<sup>2</sup> and ANDREW R. INGLIS<sup>1</sup>

<sup>1</sup>Centre for Fusion, Space and Astrophysics, Physics Department, University of Warwick, Coventry CV8 1DA, UK

<sup>2</sup> Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Moscow 119992, Russia

Quasi-periodic pulsations (QPPs) of gamma-ray, hard X-ray and microwave emission with a period of about 40 s are found in a single loop X-class solar flare on 2005 January 1 at photon energies up to 2–6 MeV with the CORONAS-F SOlar Neutrons and Gamma-rays (SONG) experiment, in hard X-ray and low energy gamma-ray channels of RHESSI and in both 17 GHz and 34 GHz channels of the Nobeyama Radioheliograph [1]. Periodogram and correlation analyses show that the 40 s QPPs of microwave, hard X-ray, and gamma-ray emission are almost synchronous in all observation bands. Analysis of the spatial structure of hard X-ray and low energy (80–225 keV) gamma-ray QPP with RHESSI reveals synchronous while asymmetric QPP at both footpoints of the flaring loop. The difference between the averaged hard X-ray fluxes coming from the two footpoint sources is found to oscillate with a period of about 13 s for five cycles in the highest emission stage of the flare. The proposed mechanism generating the 40 s QPP is a triggering of magnetic reconnection by a kink oscillation in a nearby loop [2]. The 13 s periodicity could be produced by the second harmonics of the sausage mode of the flaring loop.

## References

- [1] V.M. Nakariakov, C. Foullon, I.N. Myagkova, A.R. Inglis, Astrophys. J. 708, L47 (2010).
- [2] V.M. Nakariakov, C. Foullon, E. Verwichte, N.P. Young, Astron. Astrophys. 452, 343 (2006)