Category: OA8 Preferred Mode of Presentation: Oral

Hurricanes in the South Pacific Ocean; Characteristics and Consequences for Island Environments

JAMES P. TERRY¹ and SIMON MCGREE²

¹*Head of Geography Department, The University of the South Pacific, Suva, Fiji Islands* ²*Climate Change Unit, Fiji Meteorological Services, Nadi Airport, Fiji*

The most extreme meteorological conditions in the tropical South Pacific occur during hurricanes (tropical cyclones) and small islands are vulnerable to their effects. Hurricanes are relatively infrequent events in the south west Pacific averaging 9.4 events per season, but in optimal conditions they form quickly and bring violent winds, torrential precipitation, destructive waves, dangerous storm surge and river flooding. This paper outlines the climatology of South Pacific hurricanes. Cyclone formation and zone of origin are examined, and storm characteristics (e.g. frequency, size, structure, track movements and longevity) are described. The behaviour and impacts of recent catastrophic storms are illustrated, such as hurricane Zoe which struck the outer Solomon Islands in December 2002 and hurricane Ami which devastated many northern Fiji Islands in January 2003.

Different types of Pacific islands, e.g. volcanic, limestone and coral islands, suffer a range of environmental effects during hurricanes. If a storm makes landfall on a high volcanic island, the mountainous terrain causes orographic uplift of the rain bands, delivering exceptional and widespread precipitation. Both total amounts and intensities are extreme (e.g. 610 mm in 24 hours). The normal arrangement of rain clouds is in a spiral pattern with gaps of clear weather between heavy downpours, but vertical shearing may distort the cloud pattern, distributing moisture unevenly [1] and making localised rainfall prediction difficult. Several hurricane examples demonstrate the magnitude of storm surges and coastal inundation, shoreline erosion, destruction of coral reefs, landslides and hillslope denudation, and large river floods. All of these physical impacts cause human hardship, especially for the poor, and place difficult economic burdens on developing Pacific island states with limited resources.

Since hurricanes are often associated with the ENSO phenomenon, this link is discussed. During the developmental phase of El Niños, hurricanes affecting South Pacific islands are generated as the eastwardmigrating pool of warm ocean water passes across the north. Regarding future climate change in the South Pacific, many uncertainties remain. However, the region is likely to experience more sustained El Niñolike conditions, owing to global atmospheric and ocean warming [2, 3]. Implications for hurricane characteristics include the potential for increased intensities [4, 5], greater storm numbers [6], changes to the present spatial distribution of storm origins [7] and farther poleward travel after vortex formation [8]. Such changes will only worsen the vulnerability of island nations to hurricane impacts, so it is important for climate scientists to investigate evolving hurricane behaviour in the South Pacific, in order to advance appropriate strategies for adaptation.

References

- [1] Terry, J.P. and Raj, R. 1999 Island environment and landscape responses to 1997 tropical cyclones in Fiji. *Pacific Science*, 13, 257-272.
- [2] Trenberth, K. and Hoar, T.J. 1997 El Niño and climate change. Geophysical Research Letters 24, 3057-3060.
- [3] Timmermann, A., Oberhuber, J., Bacher, A. Esch, M., Latif, M. and Roeckner, E. 1999 Increased El Niño frequency in a climate model forced by future greenhouse warming. *Nature* 398, 694-697.
- [4] Emanuel, K.A. 1987 The dependence of hurricane intensity on climate. *Nature* 326, 483-485.
- [5] Whetton, P., Jones, R., Hennessy, K., Suppiah, R., Walsh, K. and Cai, W. 2000 Scenarios of climate change for the Pacific Islands. *Pacific Islands Climate Change Conference*, 3-7 April 2000, Rarotonga, Cook Islands. South Pacific Regional Environment Programme, Apia, Samoa.
- [6] Basher, R.E. and Zheng, X. 1995 Tropical cyclones in the southwest Pacific: spatial patterns and relationships to Southern Oscillation and sea surface temperature. *Journal of Climate* 8, 1249-1260.
- [7] Revell, C.G. and Goulter, S.W. 1986 South Pacific tropical cyclones and the Southern Oscillation. *Monthly Weather Review* 114, 1138-1145.
- [8] Walsh, K.J.E. and Katzfey, J.J. 2000 the impact of climate change on the poleward movement of tropical cyclone-like vortices in a regional climate model. *Journal of Climate* 13, 1116-1132.