

Postseismic deformations following the 2004 SE off Kii peninsula earthquake sequence and its tectonic implications

MANABU HASHIMOTO, KENSUKE ONOUE, FUMIO OHYA, YOSHINOBU HOSO,
YASUYOSHI FUJITA, KAZUTOSHI SATO and KOHEI SEGAWA

Disaster Prevention Research Institute, Kyoto University

On September 5, 2004, two earthquakes of M7.1 and 7.4 hit the central and southwest Japan. Hypocenters of these events are located SE off Kii peninsula and right beneath the axis of the Nankai trough. Aftershocks are aligned in two trends: one is parallel to the Nankai trough and the other is oblique to it. These events have almost pure thrust mechanism, but their P axes are in the NS direction different from the relative motion between the subducting Philippine Sea and overlying continental plates and nodal planes are very steep. However, some large aftershocks have nearly strike-slip fault mechanism. These strike-slip type aftershocks occurred along the oblique distribution to the Nankai trough.

We made an urgent continuous observation of denser network of GPS than GEONET in order to detect postseismic deformations following these events. Since we established this network in 2001 and repeated observations for a week or two every spring, we can compare the coordinates after the earthquakes with those in spring in 2004. We used dual frequency receivers. We adopted a standard scheme of continuous observation of 30sec sampling and elevation mask of 10 degrees. We continued observation for one and half month after the events and analyzed the data with GIPSY OASIS II. In this analysis the precise point positioning was adopted using precise ephemerides by JPL.

Coseismically most GPS stations moved south, but those on the west side of Kii peninsula moved west to southwestward. This pattern of horizontal displacements can explained by a thrust fault trending parallel to the Nankai trough. We cannot determine the dip direction on the basis of observed displacements. During the first three weeks at the postseismic stage, the E-W components increased slightly at all stations (eastward shift), then decreased. Southward movements up to 1cm are dominant even at the stations on the west side of Kii peninsula. This result suggests that the coseismic and afterslip are different from each other. One possible explanation of postseismic deformation is afterslip on the fault plane that ruptured in the main shock or its extension. Right lateral strike-slip along the oblique distribution of aftershocks may not contribute to the postseismic deformations so much, since it will produce NW-ward movements in Kii peninsula.