

## High Ductility of K-Feldspar and Development of Granitic Banded Ultramylonite in Ryoke Metamorphic Belt, SW Japan

KAZUHIKO ISHII<sup>1</sup>, KYUICHI KANAGAWA<sup>2</sup>, NORIO SHIGEMATSU<sup>3</sup>,  
TAKAMOTO OKUDAIRA<sup>4</sup>

<sup>1</sup>*Depart. Phys. Sci., Osaka Pref. Univ., Japan*

<sup>2</sup>*Depart. Earth Sci., Chiba Univ., Japan*

<sup>3</sup>*Geological Survey of Japan*

<sup>4</sup>*Depart. Geosci., Osaka City Univ., Japan*

Granitic mylonites from a greenschist facies mylonite zone in Ryoke metamorphic belt, SW Japan show three types of microstructure with increasing mylonitization: (1) mylonites, (2) banded mylonites and (3) banded ultramylonites. The banded structure is composed of quartz layers, biotite layers, K-feldspar layers, plagioclase > K-feldspar (> quartz + biotite) layers and K-feldspar > plagioclase > quartz layers. Microstructural and textural investigations by cathodoluminescence (CL), scanning electron microscope (SEM) and electron back-scatter diffraction (EBSD) techniques show that different kind of layer was deformed by different processes including dislocation creep of quartz and K-feldspar and interstitial filling the space between clasts by K-feldspar from solution. Dominant processes of the grain-size reduction were fracturing of plagioclase, myrmekite formation around K-feldspar and dynamic recrystallization of quartz. The development of the banded structure was promoted by high mobility (solution transfer) and ductility (dislocation creep) of K-feldspar.